

FIX-EULER-MAHONIAN STATISTICS ON WREATH PRODUCTS

HILARION L. M. FALIHARIMALALA AND JIANG ZENG

ABSTRACT. In 1997 Clarke et al. studied a q -analogue of Euler's difference table for $n!$ using a key bijection Ψ on symmetric groups. In this paper we extend their results to the wreath product of a cyclic group with the symmetric group. In particular we obtain a new Mahonian statistic $fmaf$ on wreath products. We also show that Foata and Han's two recent transformations on the symmetric groups provide indeed a factorization of Ψ .

Dedicated to Dennis Stanton on the occasion of his 60th birthday

1. INTRODUCTION

For positive integers $\ell, n \geq 1$, define the colored set $\Sigma_{\ell,n} = \{\zeta^j k \mid 0 \leq j \leq \ell-1, 1 \leq k \leq n\}$. Let $G_{\ell,n}$ be the set of permutations σ of $\Sigma_{\ell,n}$ such that $\sigma(\zeta^j k) = \zeta^j \sigma(k)$ for any $k \in [n]$ and $0 \leq j \leq \ell-1$. It is known that $G_{\ell,n}$ is isomorphic to the wreath product $C_\ell \wr S_n$, where S_n denotes the group of permutations of $[n] := \{1, 2, \dots, n\}$ and C_ℓ the ℓ -cyclic group generated by $\zeta = e^{2i\pi/\ell}$. In the last two decades many authors have been trying to extend various enumerative results on symmetric groups to other classical reflection groups, see [2, 3, 5, 7, 10, 6] and the references cited there. In particular Adin and Roichman [3] introduced the flag major index $fmaj$ on wreath products of a cyclic group with the symmetric group, and Haglund et al. [10] proved that

$$\sum_{\sigma \in G_{\ell,n}} q^{\text{fmaj } \sigma} = [\ell]_q [2\ell]_q \cdots [n\ell]_q, \quad (1.1)$$

where $[n]_q$ is the q -integer $1 + q + \cdots + q^{n-1}$. In this paper we shall consider some natural generalizations of (1.1) by studying a wreath product analogue of Euler's q -difference table $\{g_{\ell,n}^m(q)\}_{m \geq n \geq 0}$ defined by the following recurrence:

$$\begin{cases} g_{\ell,n}^n(q) = [\ell]_q [2\ell]_q \cdots [n\ell]_q, \\ g_{\ell,n}^m(q) = g_{\ell,n}^{m+1}(q) - q^{\ell(n-m-1)} g_{\ell,n-1}^m(q) \quad (0 \leq m \leq n-1). \end{cases} \quad (1.2)$$

For example, when $\ell = 1$ and $\ell = 2$, the first values of $g_{\ell,n}^m(q)$ are given as follows:

- $\ell = 1$

$n \backslash m$	0	1	2	3
0	1			
1	0	1		
2	q	q	$1+q$	
3	$q[2]_q$	$q[3]_q$	$q^3 + 2q^2 + q$	$(1+q)(1+q+q^2)$

Date: December 1, 2009.

• $\ell = 2$

$n \backslash m$	0	1	2	3
0	1			
1	q	$[2]_q$		
2	$q[4]_q + q^2$	$q[4]_q + q^2[2]_q$	$[2]_q[4]_q$	
3	$g_{2,3}^0$	$g_{2,3}^1$	$g_{2,3}^2$	$[2]_q[4]_q[6]_q$

where

$$\begin{aligned} g_{2,3}^2 &= q^9 + 3q^7 + 5q^7 + 7q^6 + 8q^5 + 7q^4 + 5q^3 + 3q^2 + q, \\ g_{2,3}^1 &= q^9 + 3q^8 + 5q^7 + 6q^6 + 6q^5 + 5q^4 + 4q^3 + 3q^2 + q, \\ g_{2,3}^0 &= q^9 + 2q^8 + 4q^7 + 4q^6 + 5q^5 + 5q^4 + 4q^3 + 3q^2 + q. \end{aligned}$$

It is remarkable that $g_{\ell,n}^m(q)$ are polynomials in q with non-negative integral coefficients. For $\ell = 1$, Clarke et al. [4] proved that the entry $g_{1,n}^m(q)$ is actually the generating function for a subset of S_n by the Mahonian statistic *maf*. Their proof is based on a key bijection Ψ on S_n transforming the statistic *maf* to the statistic *major index*. Our first aim is to show that the results in [4] can be readily extended to wreath products. More precisely, we will find a combinatorial interpretation and an explicit formula for $g_{\ell,n}^m(q)$ by introducing a new Mahonian statistic *fmaf* on the wreath products and by extending Clarke et al.'s bijection Ψ to the colored permutations. On the other hand, Foata and Han [7] have recently constructed two new transformations on symmetric groups and *noticed* that the composition of their two transformations has some properties in common with Clarke et al.'s bijection Ψ in [4]. A natural question is then to ask wheather these two algorithms are *identical*. Our second aim of this paper is to settle this open question.

2. DEFINITIONS AND MAIN RESULTS

For $x = \zeta^j k \in \Sigma_{\ell,n}$, we write $x = \varepsilon_x |x|$ with

$$\varepsilon_x = \zeta^j \in C_\ell \quad \text{and} \quad |x| = k \in [n].$$

If ℓ is small, it is convenient to write j bars over i instead of $\zeta^j i$, thus $\zeta^2 4 = \bar{\bar{4}}$. By ordering the elements of C_ℓ as $1 > \zeta > \zeta^2 > \dots > \zeta^{\ell-1}$, we can define a linear order on the alphabet $\Sigma_{\ell,n}$ as follows:

$$\varepsilon_{x_1} |x_1| < \varepsilon_{x_2} |x_2| \Leftrightarrow [\varepsilon_{x_1} < \varepsilon_{x_2}] \text{ or } [(\varepsilon_{x_1} = \varepsilon_{x_2}) \text{ and } (|x_1| < |x_2|)]. \quad (2.1)$$

For example, the elements of $\Sigma_{4,4}$ are ordered as follows:

$$\bar{\bar{1}} < \bar{\bar{2}} < \bar{\bar{3}} < \bar{\bar{4}} < \bar{1} < \bar{2} < \bar{3} < \bar{4} < 1 < 2 < 3 < 4.$$

Recall that $i \in [n]$ is a *fixed point* of $\sigma \in G_{\ell,n}$ if $\sigma(i) = i$. Let $\text{FIX}(\sigma)$ be the set of fixed points of σ and $\text{fix } \sigma$ the cardinality of $\text{FIX}(\sigma)$. The colored permutation σ has a *descent* at $i \in \{1, 2, \dots, n-1\}$ if $\sigma(i) > \sigma(i+1)$ and i is called a *descent place* of σ . Let $\text{DES}(\sigma)$ be the set of descent places of σ . The *major index* of σ , denoted by $\text{maj } \sigma$, is the sum of all the descent places of σ . If $y_1 y_2 \dots y_m$ is the word obtained by deleting the fixed points of σ and

$y_i = \varepsilon_i |y_i|$ ($1 \leq i \leq m$), writing $z_i = \varepsilon_i \text{rank}(|y_i|)$ with ‘rank’ being the increasing bijection from $\{|y_1|, |y_2|, \dots, |y_m|\}$ to $[m]$, then the *derangement part* of σ is defined to be

$$\text{Der}(\sigma) = z_1 z_2 \cdots z_m. \quad (2.2)$$

We now define the main Eulerian and Mahonian statistics on the wreath product $G_{\ell,n}$. If $\sigma = x_1 \dots x_n \in G_{\ell,n}$ then the statistics des_ℓ , exc and col are defined by

$$\text{des}_\ell \sigma = \sum_{i=1}^{n-1} \chi(x_i > x_{i+1}), \quad \text{exc} \sigma = \sum_{i=1}^n \chi(x_i > i), \quad \text{col} \sigma = \sum_{j=0}^{\ell-1} j \cdot |\text{COL}_j(\sigma)|,$$

where $\text{COL}_j(\sigma) = \{i \in [n] : \frac{x_i}{|x_i|} = \zeta^j\}$ and $\chi(A) = 1$ if A is true and 0 otherwise. When $\ell = 1$ we shall write $\text{des} = \text{des}_\ell$. The maj and maf statistics are defined by

$$\text{maj} \sigma = \sum_{i=1}^{n-1} i \cdot \chi(x_i > x_{i+1}), \quad \text{and} \quad \text{maf} \sigma = \text{maj} \text{Der}(\sigma) + \sum_{j=1}^k (i_j - j),$$

where $\text{FIX}(\sigma) = \{i_1, i_2, \dots, i_k\}$. The *flag-maj* fmaj and *flag-maf* fmaf statistic are defined by

$$\text{fmaj} \sigma = \ell \cdot \text{maj} \sigma + \text{col}(\sigma), \quad \text{and} \quad \text{fmaf} \sigma = \ell \cdot \text{maf} \sigma + \text{col}(\sigma).$$

Remark. While the statistic fmaj was first introduced by Adin and Roichman [3], the statistic fmaf is new and reduces to maf of Clarke et al. [4] for $\ell = 1$.

Example. If $\sigma = \mathbf{1\bar{8}346\bar{2}7\bar{5}9} \in G_{4,9}$, then $\text{maj} \sigma = 1 + 5 + 7 = 13$, $\text{FIX} \sigma = \{1, 3, 4, 7, 9\}$ and $\text{Der}(\sigma) = \mathbf{4\bar{3}\bar{1}\bar{2}}$. Therefore $\text{maj} \text{Der}(\sigma) = 2$ and

$$\text{maf} \sigma = 2 + ((1 - 1) + (3 - 2) + (4 - 3) + (7 - 4) + (9 - 5)) = 11.$$

Since $\text{COL}_0(\sigma) = \{1, 3, 4, 5, 7, 9\}$, $\text{COL}_1(\sigma) = \{2, 8\}$, $\text{COL}_2(\sigma) = \{6\}$, $\text{COL}_3(\sigma) = \emptyset$, we have $\text{col}(\sigma) = 0 \times 6 + 1 \times 2 + 2 \times 1 = 4$. Finally,

$$\text{fmaj} \sigma = 4 \times 13 + 4 = 56, \quad \text{and} \quad \text{fmaf} \sigma = 4 \times 11 + 4 = 48.$$

Remark. Other notions of descents have also been considered previously, see [13]. For the length function of the wreath product $C_\ell \wr S_n$ we refer the reader to [9].

We first show that the statistics $(\text{fix}, \text{fmaf})$ and $(\text{fix}, \text{fmaj})$ are equidistributed on $G_{\ell,n}$ and their common distribution has an explicit formula.

Theorem 1. *The triple statistics $(\text{fmaf}, \text{exc}, \text{fix})$ and $(\text{fmaj}, \text{exc}, \text{fix})$ are equidistributed on $G_{\ell,n}$. Moreover the common generating function*

$$\sum_{\sigma \in G_{\ell,n}} q^{\text{fmaf} \sigma} x^{\text{fix} \sigma} = \sum_{\sigma \in G_{\ell,n}} q^{\text{fmaj} \sigma} x^{\text{fix} \sigma} \quad (2.3)$$

has the explicit formula

$$g_{\ell,n}(q, x) := [\ell]_q [2\ell]_q \cdots [n\ell]_q \sum_{k=0}^n \frac{(x-1)(x-q^\ell) \cdots (x-q^{\ell(k-1)})}{[\ell]_q [2\ell]_q \cdots [k\ell]_q}. \quad (2.4)$$

Remark. When $\ell = 1$ Gessel and Reutenauer [8] first proved that the generating function of $(\text{fmaj}, \text{fix})$ on S_n is given by (2.4). For general ℓ and $x = 1$ we recover Haglund et al's formula (1.1) for the generating function of fmaj on $G_{\ell,n}$. For $x = 0$, we derive from (2.3) and (2.4) an explicit formula for the colored q -derangement number:

$$d_{\ell,n}(q) := \sum_{\sigma \in \mathcal{D}_{\ell,n}} q^{\text{fmaj } \sigma} = [\ell]_q [2\ell]_q \cdots [n\ell]_q \sum_{k=0}^n \frac{(-1)^k q^{\ell \binom{k}{2}}}{[\ell]_q [2\ell]_q \cdots [k\ell]_q}. \quad (2.5)$$

The $\ell = 1$ and $\ell = 2$ cases of (2.5) were first obtained by Gessel (unpublished) and Wachs [14], and Chow [5], respectively. Finally (2.5) yields immediately the following recurrence relation for the colored q -derangement numbers:

$$d_{\ell,n+1}(q) = [\ell n + \ell]_q d_{\ell,n}(q) + (-1)^{n+1} q^{\ell \binom{n+1}{2}}. \quad (2.6)$$

Introduce the q -shifted factorials

$$(a; q)_0 = 1, \quad (a; q)_n = \prod_{k=0}^{n-1} (1 - aq^k), \quad n = 1, 2, \dots, \text{ or } \infty,$$

then the q -binomial coefficients are defined by

$$\begin{bmatrix} n \\ m \end{bmatrix}_q = \frac{(q; q)_n}{(q; q)_m (q; q)_{n-m}}, \quad n \geq m \geq 0.$$

Instead of the colored q -Euler table (1.2), as Clarke et al. [4], we can consider a more general triangle than (1.2) by taking $g_{\ell,n}^n(q, x) := g_{\ell,n}(q, x)$ as the diagonal coefficients and replace the recurrence relation in (1.2) by

$$g_{\ell,n}^m(q, x) = g_{\ell,n}^{m+1}(q, x) - xq^{\ell(n-m-1)} g_{\ell,n-1}^m(q, x) \quad (0 \leq m \leq n-1). \quad (2.7)$$

For $n \geq m \geq 0$, denote by $G_{\ell,n}^m$ the set of permutations σ in $G_{\ell,n}$ such that $\text{FIX}(\sigma) \subset \{n-m+1, \dots, n-1, n\}$, i.e., $\sigma(i) \neq i$ for $i \leq n-m$. In particular we have $G_{\ell,n}^n = G_{\ell,n}$ and $\mathcal{D}_{\ell,n} := G_{\ell,n}^0$ is the set of colored derangements of order n . The following theorem gives a full description of $g_{\ell,n}^m(q, x)$, which generalizes the previous results in [4] for $\ell = 1$ and [6] for $q = 1$, respectively.


Theorem 2. For $n \geq m \geq 0$ we have the following explicit formula:

$$g_{\ell,n}^m(q, x) = \sum_{k=0}^{n-m} (-x)^k \begin{bmatrix} n-m \\ k \end{bmatrix}_{q^\ell} q^{\ell \binom{k}{2}} g_{\ell,n-k}(q, x) \quad (2.8)$$

and the combinatorial interpretation:

$$g_{\ell,n}^m(q, x) = \sum_{\sigma \in G_{\ell,n}^m} q^{\text{fmaj } \sigma} x^{\text{fix } \sigma}. \quad (2.9)$$

Remark. The two statistics fmaj and fmaj are identical on $\mathcal{D}_{\ell,n}$, equidistributed on $G_{\ell,n}$, but not equidistributed on the set $G_{\ell,n}^m$ for $0 < m < n$.

FIGURE 1. The Foata-Han factorization of Clarke et al.'s bijection Ψ

Let $0 \leq m \leq n$ and v be a nonempty word of length m on the alphabet $[n]$. Denote by $\text{Sh}(0^{n-m}v)$ the set of all *shuffles* of the words 0^{n-m} and v , that is, the set of all words it is possible to construct using $(n-m)$ 0's and the letters in v by preserving the order of all the letters in v . For any word $w = x_1 \dots x_n$ in $\text{Sh}(0^{n-m}v)$, we call *pillar* any non zero letter of w and write $\text{Pil}(w) := v$ and $\text{Zero } w := \{i : 1 \leq i \leq n, x_i = 0\}$. Let $\text{pil } w$ be the length of $\text{Pil } w$ and $\text{zero } w$ be the cardinality of $\text{Zero } w$. Clearly w is completely characterized by the pair $(\text{Zero } w, \text{Pil } w)$. Given a permutation $\sigma = \sigma(1)\sigma(2) \dots \sigma(n+m) \in S_{n+m}$, let $(j_1, j_2, \dots, j_m)$ be the increasing sequence of the integers k such that $\sigma(k) \neq k$ for $1 \leq k \leq n$. The word $w = x_1 x_2 \dots x_{n+m}$ derived from σ by replacing each fixed point by 0 and each other letter $\sigma(j_k)$ by $\text{rank}(\sigma(j_k))$ will be denoted by $\text{ZDer}(\sigma)$. For example, if $\sigma = \mathbf{1\ 8\ 3\ 4\ 6\ 2\ 7\ 5\ 9} \in S_9$ then $\text{ZDer}(\sigma) = 0\ 4\ 0\ 0\ 3\ 1\ 0\ 2\ 0$. Let

$$\mathcal{S}_n^{\text{Der}} := \bigcup_{m,v} \text{Sh}(0^{n-m}v) \quad (0 \leq m \leq n, v \in \mathcal{D}_m).$$

It is obvious (see [7, Proposition 1.3]) that the map ZDer is a bijection from S_n to $\mathcal{S}_n^{\text{Der}}$. Recently Foata and Han [7] have constructed two transformations Φ and $\mathbf{F} := \mathbf{F}_3$ on $\mathcal{S}_n^{\text{Der}}$ (see Section 5), apparently related to Ψ . Our third object is to show that their two bijections provide indeed a factorization of Clarke et al.'s bijection Ψ .

Theorem 3. *We have*

$$\Psi = \text{ZDer}^{-1} \circ \mathbf{F} \circ \Phi^{-1} \circ \text{ZDer}. \quad (2.10)$$

In other words, the diagram in Figure 1 is commutative.

The rest of this paper is organized as follows. In Section 3, after recalling Clarke et al.'s bijection Ψ , we prove Theorem 1. The proof of Theorem 2 is given in Section 4. In Section 5, after recalling Foata and Han's two transformations Φ and $\mathbf{F}$, we prove Theorem 3 and postpone the proof of three technical lemmas to Section 6.

3. PROOF OF THEOREM 1

We first recall Clarke et al.'s bijection Ψ , which will also be used in Section 5, and then show that one can extend Ψ to $G_{\ell,n}$ with the following property:

$$(\text{maf}, \text{exc}, \text{fix}, \text{col})\Psi(\sigma) = (\text{maj}, \text{exc}, \text{fix}, \text{col})\sigma, \quad \forall \sigma \in G_{\ell,n}. \quad (3.1)$$

3.1. Clarke et al.'s bijection Ψ . Let $\sigma = x_1 x_2 \dots x_n \in S_n$ with $x_0 = x_{n+1} = +\infty$. For $0 \leq i \leq n$, a pair $(i, i+1)$ of positions is the j -th slot of σ provided that $x_i \neq i$ and that σ has $i-j$ fixed points f such that $f < i$. Of course, the j -th slot is $(j, j+1)$ if σ is a derangement. Clearly we can insert a fixed point into the j -th slot and obtain the permutation

$$\langle \sigma, j \rangle = x'_1 x'_2 \dots x'_i (i+1) x'_{i+1} \dots x'_n, \quad (3.2)$$

where $x' = x$ if $x \leq i$ and $x' = x+1$ if $x > i$ for all $x \in [n]$.

Now, if σ is a derangement in S_n and $(i_1, i_2, \dots, i_m)$ a sequence of integers such that $0 \leq i_1 \leq i_2 \leq \dots \leq i_m \leq n$, we can insert successively m fixed points in σ and obtain a permutation τ in S_{n+m} :

$$\tau = \langle \sigma, i_1, \dots, i_m \rangle = \langle \langle \sigma, i_1, \dots, i_{m-1} \rangle, i_m \rangle. \quad (3.3)$$

Note that $\text{Der}(\tau) = \sigma$ and the fixed points of the last permutation are $i_1+1, i_2+2, \dots, i_m+m$. Conversely, any permutation $\tau \in S_{m+n}$ with m fixed points can be written as (3.3) in one and only one way. Thus, if $S(\sigma, m)$ denotes the set of permutations in S_{n+m} with derangement part $\sigma \in \mathcal{D}_n$, then

$$S(\sigma, m) = \{ \langle \sigma, i_1, \dots, i_m \rangle \mid 0 \leq i_1 \leq i_2 \leq \dots \leq i_m \leq n \}.$$

Let $\sigma \in S_n$. The j -th slot $(i, i+1)$ of σ is said to be *green* if $\text{des}\langle \sigma, j \rangle = \text{des}\sigma$, *red* if $\text{des}\langle \sigma, j \rangle = \text{des}\sigma + 1$. We assign *values* from 0 to g to the green slots of σ from right to left, and values from $g+1$ to n to the red slots from left to right. Denote the value of the j -th slot by g_j . The bijection $\Psi : S(\sigma, m) \rightarrow S(\sigma, m)$ is defined by induction on $m \geq 0$ as follows:

- (1) Ψ is the identity mapping on $S(\sigma, 0)$.
- (2) Define Ψ on $S(\sigma, 1)$ by

$$\Psi \langle \sigma, i \rangle = \langle \sigma, g_i \rangle. \quad (3.4)$$

- (3) Let $m > 1$ and suppose that Ψ has been defined on $S(\sigma, k)$ for $0 \leq k \leq m-1$. Consider $\tau = \langle \sigma, i_1, \dots, i_m \rangle$. Suppose that the i_m -th slot of σ is green. Then, if $\Psi \langle \sigma, i_1, \dots, i_{m-1} \rangle = \langle \sigma, j_2, \dots, j_m \rangle$, we define

$$\Psi(\tau) = \langle \sigma, g_{i_m}, j_2, \dots, j_m \rangle. \quad (3.5)$$

Suppose that the i_m -th slot of σ is red. Let k be the smallest positive integer such that $i_{m-k} < i_m$. Thus $i_{m-k} < i_{m-k+1} = \dots = i_m$. Then,

- if $k = m$ we define

$$\Psi(\tau) = \langle \sigma, \underbrace{g_{i_m} - i_m, \dots, g_{i_m} - i_m}_{m-1 \text{ terms}}, g_{i_m} \rangle. \quad (3.6)$$

- if $k < m$ and $\Psi \langle \sigma, i_1, \dots, i_{m-k} \rangle = \langle \sigma, j_1, \dots, j_{m-k} \rangle$, we define

$$\Psi(\tau) = \langle \sigma, \underbrace{g_{i_m} - i_m, \dots, g_{i_m} - i_m}_{k-1 \text{ terms}}, j_1 + 1, \dots, j_{m-k} + 1, g_{i_m} \rangle. \quad (3.7)$$

3.2. Generalization of Ψ to $G_{\ell,n}$. To extend the insertion algorithm (3.3) to colored permutations we just need to modify equation (3.2) as follows:

$$\langle \sigma, j \rangle = x'_1 x'_2 \dots x'_i (i+1) x'_{i+1} \dots x'_n, \quad (3.8)$$

where $x' = x$ if $|x| \leq i$ and

$$x' = x + 1 := \varepsilon_x(|x| + 1), \quad (3.9)$$

if $|x| > i$ for all $x \in \Sigma_{\ell,n}$. Thus each colored permutation $\tau \in G_{\ell,n}$ can be written as (3.3).

Example. Let $\tau = 1 \bar{9} 3 10 5 6 7 4 2 \bar{8} \in G_{3,10}$. Then $\sigma := \text{Der}(\tau) = \bar{4} 5 2 1 \bar{3} \in \mathcal{D}_{3,5}$ and $\tau = \langle \sigma, 0, 1, 2, 2, 2 \rangle$. Indeed,

$$\begin{aligned} \langle \sigma, 0 \rangle &= 1 \bar{5} 6 3 2 \bar{4}, \\ \langle \sigma, 0, 1 \rangle &= 1 \bar{6} 3 7 4 2 \bar{5}, \\ \langle \sigma, 0, 1, 2 \rangle &= 1 \bar{7} 3 8 5 4 2 \bar{6}, \\ \langle \sigma, 0, 1, 2, 2 \rangle &= 1 \bar{8} 3 9 5 6 4 2 \bar{7}, \\ \langle \sigma, 0, 1, 2, 2, 2 \rangle &= 1 \bar{9} 3 10 5 6 7 4 2 \bar{8}. \end{aligned}$$

So $\tau = \langle \sigma, 0, 1, 2, 2, 2 \rangle$.

As in the symmetric group we say that the i -th slot ($0 \leq i \leq n$) of $\sigma \in G_{\ell,n}$ is green if $\text{des}_{\ell} \langle \sigma, i \rangle = \text{des}_{\ell} \sigma$ and red if $\text{des}_{\ell} \langle \sigma, i \rangle = \text{des}_{\ell} \sigma + 1$. In the same way, we can assign a value to each slot of σ . Now, let σ be a derangement and $G(\sigma, m) := \{\tau \in G_{\ell,n+m} \mid \text{Der}(\tau) = \sigma\}$. We can extend Ψ to a bijection on $G(\sigma, m)$. It follows that $\text{Der}(\Psi(\tau)) = \text{Der}(\tau)$ for any $\tau \in G(\sigma, m)$. Since $\text{exc } \tau = \text{exc}(\text{Der}(\tau))$ and $\text{col } \tau = \text{col}(\text{Der}(\tau))$, we have immediately

$$(\text{exc}, \text{fix}, \text{col})\Psi(\tau) = (\text{exc}, \text{fix}, \text{col})\tau.$$

It remains to verify $\text{maj } \tau = \text{maf}(\Psi(\tau))$. On symmetric groups, the proof of the latter equality is based on the following result.

Lemma 1 (Clarke et al.). *Let σ be a derangement in symmetric group and g_i the value of its i -th slot then*

$$\text{maj} \langle \sigma, i \rangle = \text{maj } \sigma + g_i. \quad (3.10)$$

Now, the substitution $x \rightarrow x'$ in (3.8) is compatible with the linear order (see (2.1)) on the alphabet $\Sigma_{\ell,n}$, namely

$$\forall a, b \in \Sigma_{\ell,n} \quad a < b \iff a' < b',$$

hence $\text{DES}(x'_1 x'_2 \dots x'_i) = \text{DES}(x_1 x_2 \dots x_i)$ and $\text{DES}(x'_{i+1} \dots x'_n) = \text{DES}(x_{i+1} \dots x_n)$. So the proof of Lemma 1 in [4] remains valid when we replace a derangement σ by a any colored derangement. Then we derive (3.1) as in [4].

Example. Let $\tau = 1 \bar{9} 3 10 5 6 7 4 2 \bar{8} \in G_{3,10}$. We have $\text{DES } \tau = \{1, 4, 7, 8, 9\}$ and $\text{maj } \tau = 29$, $\text{Der}(\tau) = \sigma = \bar{4} 5 2 1 \bar{3}$; $\text{maj } \sigma = 9$. By the previous example, we can write $\tau = \langle \sigma, 0, 1, 2, 2, 2 \rangle$. Note that slots 1,3,4,5 are *green*, while slots 0 and 2 are *red*. So the sequence of values of the slots is $(g_0, g_1, g_2, g_3, g_4, g_5) = (4, 3, 5, 2, 1, 0)$. The algorithm Ψ goes as follows:

- $\Psi\langle\sigma, 0\rangle = \langle\sigma, g_0\rangle = \langle\sigma, 4\rangle$;
- Since 1 is green, $\Psi\langle\sigma, 0, 1\rangle = \langle\sigma, g_1, 4\rangle = \langle\sigma, 3, 4\rangle$;
- Since 2 is red, $\Psi\langle\sigma, 0, 1, 2, 2, 2\rangle = \langle\sigma, g_2 - 2, g_2 - 2, 3 + 1, 4 + 1, g_2\rangle = \langle\sigma, 3, 3, 4, 5, 5\rangle$.

Thus $\Psi(\tau) = \bar{6} \ 8 \ 2 \ 4 \ 5 \ 1 \ 7 \ \bar{3} \ 9 \ 10$. Note that

$$\begin{aligned} \text{maf } \Psi(\tau) &= \text{maj } \sigma + 3 + 3 + 4 + 5 + 5 = 29 = \text{maj}(\tau), \\ \text{fmaf } \Psi(\tau) &= \text{fmaj}(\tau) = 3 \times 29 + 3 = 90. \end{aligned}$$

3.3. Proof of (2.4). Recall Cauchy's q -binomial formula (cf. [1, p.17]):

$$\sum_{n \geq 0} \frac{(a; q)_n}{(q; q)_n} u^n = \frac{(au; q)_\infty}{(u; q)_\infty}.$$

In particular we have Euler's formula by taking $a = 0$:

$$e_q(u) := \sum_{n \geq 0} \frac{u^n}{(q; q)_n} = \frac{1}{(u; q)_\infty}. \quad (3.11)$$

Let $f_{\ell, n}(q, x) = \sum_{\sigma \in G_{\ell, n}} x^{\text{fix } \sigma} q^{\text{fmaf } \sigma}$. By the insertion algorithm, we can write any permutation $\sigma \in G_{\ell, n}$ with $n - k$ ($0 \leq k \leq n$) fixed points as $\sigma = \langle \pi, i_1, \dots, i_{n-k} \rangle$, where $\pi \in \mathcal{D}_{\ell, k}$ and $0 \leq i_1 \leq \dots \leq i_{n-k} \leq k$. Since $\text{fmaf } \sigma = \text{fmaj } \pi + \ell(i_1 + \dots + i_{n-k})$, we have

$$\begin{aligned} f_{\ell, n}(q, x) &= \sum_{k=0}^n x^{n-k} \sum_{\pi \in \mathcal{D}_{\ell, k}} \sum_{\substack{\sigma \in G_{\ell, n} \\ \text{Der}(\sigma) = \pi}} q^{\text{fmaf } \sigma} \\ &= \sum_{k=0}^n x^{n-k} \sum_{\pi \in \mathcal{D}_{\ell, k}} q^{\text{fmaj } \pi} \sum_{0 \leq i_1 \leq \dots \leq i_{n-k} \leq k} q^{\ell(i_1 + \dots + i_{n-k})} \\ &= \sum_{k=0}^n x^{n-k} \begin{bmatrix} n \\ k \end{bmatrix}_{q^\ell} d_{\ell, k}(q). \end{aligned}$$

Therefore

$$\sum_{n \geq 0} f_{\ell, n}(q, x) \frac{u^n}{(q^\ell; q^\ell)_n} = e_{q^\ell}(xu) \sum_{k=0}^{\infty} d_{\ell, k}(q) \frac{u^k}{(q^\ell; q^\ell)_k}. \quad (3.12)$$

Since $f_{\ell, n}(q, 1) = (1 - q)^{-n} (q^\ell; q^\ell)_n$ by (1.1) and (2.3), setting $x = 1$ in (3.12) yields then

$$\sum_{n \geq 0} d_{\ell, n}(q) \frac{u^n}{(q^\ell; q^\ell)_n} = \frac{1}{1 - u/(1 - q)} \frac{1}{e_{q^\ell}(u)}.$$

Substituting this back to (3.12) we obtain

$$\sum_{n \geq 0} f_{\ell, n}(q, x) \frac{u^n}{(q^\ell; q^\ell)_n} = \frac{1}{1 - u/(1 - q)} \frac{e_{q^\ell}(xu)}{e_{q^\ell}(u)}. \quad (3.13)$$

On the other hand, by (2.4) we have

$$\begin{aligned}
\sum_{n \geq 0} g_{\ell,n}(q, x) \frac{u^n}{(q^\ell, q^\ell)_n} &= \sum_{n \geq 0} \sum_{i \geq 0} \frac{(x-1)(x-q^\ell) \cdots (x-q^{\ell(i-1)})}{(q^\ell; q^\ell)_i} \frac{u^n}{(1-q)^{n-i}} \\
&= \sum_{i \geq 0} \frac{(x^{-1}; q^\ell)_i}{(q^\ell; q^\ell)_i} (xu)^i \sum_{n \geq 0} \left(\frac{u}{1-q} \right)^n \\
&= \frac{1}{1-u/(1-q)} \frac{(u; q^\ell)_\infty}{(xu; q^\ell)_\infty},
\end{aligned}$$

which is equal to the right-hand side of (3.13) by Euler's formula (3.11). It follows that $f_{\ell,n}(q, x) = g_{\ell,n}(q, x)$.

Remark. As $(1-u)e_q(u) = e_q(qu)$, we can also write (3.13) as

$$\sum_{n \geq 0} f_{\ell,n}(q, x) \frac{u^n}{(q^\ell; q^\ell)_n} = \frac{(1-q)e_{q^\ell}(xu)}{e_{q^\ell}(q^\ell u) - qe_{q^\ell}(u)}.$$

4. PROOF OF THEOREM 2

To derive an explicit formula for $g_{\ell,n}^m(q)$ we give a more general formula, which is a variant of a result in [15, Th. 3] and may be also interesting in its own right.

Lemma 2. *Let $(a_{n,m})_{0 \leq m \leq n}$ be an array defined by*

$$\begin{cases} a_{0,m} = x_m & (m = n); \\ a_{n,m} = z_m a_{n-1,m+1} + y_n a_{n-1,m} & (0 \leq m \leq n-1). \end{cases} \quad (4.1)$$

If $e_i(y_1, y_2, \dots, y_n)$ denotes the i -th elementary symmetric polynomial of $y_1, \dots, y_n$, then

$$a_{n,m} = \sum_{k=0}^n x_{m+k} (z_m z_{m+1} \cdots z_{m+k-1}) e_{n-k}(y_1, y_2, \dots, y_n). \quad (4.2)$$

Proof. The formula is obviously true for $n = 0$ and $n = 1$. Suppose that it is true until $n - 1$. Since $(1 + y_1 t)(1 + y_2 t) \cdots (1 + y_n t) = \sum_{i=0}^n e_i(y_1, \dots, y_n) t^i$, we then have

$$\begin{aligned}
a_{n,m} &= y_n a_{n-1,m} + z_m a_{n-1,m+1} \\
&= y_n \sum_{k=0}^{n-1} (z_m z_{m+1} \cdots z_{m+k-1}) e_{n-1-k}(y_1, y_2, \dots, y_{n-1}) x_{m+k} \\
&\quad + z_m \sum_{k=0}^{n-1} (z_{m+1} z_{m+2} \cdots z_{m+k}) e_{n-1-k}(y_1, y_2, \dots, y_{n-1}) x_{m+k+1} \\
&= y_n e_{n-1}(y_1, y_2, \dots, y_{n-1}) x_m \\
&\quad + \sum_{k=0}^{n-1} \left(\prod_{j=0}^k z_{m+j} \right) (y_n e_{n-2-k}(y_1, y_2, \dots, y_{n-1}) + e_{n-1-k}(y_1, y_2, \dots, y_{n-1})) x_{m+1+k} \\
&= e_n(y_1, y_2, \dots, y_n) x_m + \sum_{k=1}^n \left(\prod_{j=0}^{k-1} z_{m+j} \right) e_{n-k}(y_1, y_2, \dots, y_n) x_{m+k}.
\end{aligned}$$

This completes the proof. $\square$

Now, specializing the array (4.1) with $x_m = g_{\ell,m}^m(q, x)$, $z_m = 1$ and $y_n = -xq^{\ell(n-1)}$, then the q -binomial formula $(1+t)(1+qt) \cdots (1+q^{n-1}t) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q q^{\binom{k}{2}} t^k$ implies that

$$e_k(1, q, q^2, \dots, q^{n-1}) = \begin{bmatrix} n \\ k \end{bmatrix}_q q^{\binom{k}{2}} \quad (0 \leq k \leq n).$$

Applying (4.2) we get

$$a_{n,m} = g_{\ell,n+m}^m(q, x) = \sum_{k=0}^n (-x)^{n-k} \begin{bmatrix} n \\ k \end{bmatrix}_{q^\ell} q^{\ell \binom{n-k}{2}} g_{\ell,m+k}^{m+k}(q, x).$$

Shifting n by $n - m$ and then replacing k by $n - m - k$ yields (2.8).

Let $f_{\ell,n}^m(q, x) := \sum q^{\text{fmaf } \sigma} x^{\text{fix } \sigma}$ ($\sigma \in G_{\ell,n}^m$) be the right-hand side of (2.9). Then $f_{\ell,n}^n(q, x) = g_{\ell,n}^n(q, x)$ by Theorem 1. For each fixed n we will show by induction on m ($0 \leq m \leq n$) that $\{f_{\ell,n}^m(q, x)\}$ satisfies the recurrence relation (1.2). For $0 \leq m \leq n - 1$ define

$$E := G_{\ell,n}^{m+1} \setminus G_{\ell,n}^m = \{\sigma \in G_{\ell,n}^m : \sigma(n - m) = n - m\}.$$

By (3.3), each permutation $\sigma \in E$ can be written as

$$\sigma = \langle \text{Der } \sigma, i_1 - 1, i_2 - 2, \dots, i_r - r \rangle,$$

where $i_1, \dots, i_r$ are the fixed points of σ arranged in increasing order ($i_1 = n - m$). Let $\sigma' = \langle \text{Der } \sigma, i_2 - 2, \dots, i_r - r \rangle$. Then the mapping $\sigma \mapsto \sigma'$ is a bijection from E to $G_{\ell,n-1}^m$ such that $\text{fix } \sigma = \text{fix } \sigma' + 1$ and $\text{fmaf } \sigma = \ell((i_1 - 1) + (i_2 - 2) + \cdots + (i_r - r)) + \text{fmaf } \text{Der } \sigma$. It

follows that $\text{fmaf } \sigma = \text{fmaf } \sigma' + \ell(n - m - 1)$. Hence

$$\begin{aligned} f_{\ell,n}^{m+1}(q, x) &= \sum_{\sigma \in G_{\ell,n}^m} q^{\text{fmaf } \sigma} x^{\text{fix } \sigma} + \sum_{\sigma' \in G_{\ell,n-1}^m} q^{\text{fmaf } \sigma' + \ell(n-m-1)} x^{\text{fix } \sigma} \\ &= f_{\ell,n}^m(q, x) + xq^{\ell(n-m-1)} f_{\ell,n-1}^m(q, x). \end{aligned}$$

This completes the proof of (2.9).

5. PROOF OF THEOREM 3

5.1. Foata-Han's first transformation Φ . Let v be a derangement of order m and $w = x_1 x_2 \cdots x_n \in \text{Sh}(0^{n-m}v)$ ($0 \leq n \leq m$). Thus $v = x_{j_1} x_{j_2} \cdots x_{j_m}$, where $1 \leq j_1 < j_2 < \cdots < j_m \leq n$. Recall that “rank” is the increasing bijection of $\{j_1, j_2, \dots, j_m\}$ onto the interval $[m]$. A positive letter x_k of w is said to be *excedent* (resp. *subexcedent*) if $x_k > \text{rank}(k)$ (resp. $x_k < \text{rank}(k)$). Accordingly, a letter is *non-subexcedent* if it is either equal to 0 or *excedent*.

We define n bijections ϕ_l ($1 \leq l \leq n$) from $\text{Sh}(0^{n-m}v)$ ($0 \leq n \leq m$) onto itself in the following manner: for each l such that $n - m + 1 \leq l \leq n$ let $\phi_l(w) := w$. When $1 \leq l \leq n - m$, let x_j denote the l -th letter of w , equal to 0, when w is read *from left to right*. Three cases are next considered (by convention, $x_0 = x_{n+1} = +\infty$):

- (1) x_{j-1}, x_{j+1} both non-subexcedent;
- (2) x_{j-1} non-subexcedent, x_{j+1} subexcedent; or x_{j-1}, x_{j+1} both subexcedent with $x_{j-1} > x_{j+1}$;
- (3) x_{j-1} subexcedent, x_{j+1} non-subexcedent; or x_{j-1}, x_{j+1} both subexcedent with $x_{j-1} < x_{j+1}$.

When case (1) holds, let $\phi_l(w) := w$. When case (2) holds, determine the *greatest* integer $k \geq j+1$ such that $x_{j+1} < x_{j+2} < \cdots < x_k < \text{rank}(k)$ so that $w = x_1 \cdots x_{j-1} 0 x_{j+1} \cdots x_k x_{k+1} \cdots x_n$ and define: $\phi_l(w) := x_1 \cdots x_{j-1} x_{j+1} \cdots x_k 0 x_{k+1} \cdots x_n$. When case (3) holds, determine the *smallest* integer $i \leq j-1$ such that $\text{rank}(i) > x_i > x_{i+1} > \cdots > x_{j-1}$ so that $w = x_1 \cdots x_{i-1} x_i \cdots x_{j-1} 0 x_{j+1} \cdots x_n$ and define $\phi_l(w) := x_1 \cdots x_{i-1} 0 x_i \cdots x_{j-1} x_{j+1} \cdots x_n$.

It is important to note that ϕ_l has no action on the 0's other than the l -th one. The mapping Φ is defined to be the composition product $\Phi := \phi_1 \phi_2 \cdots \phi_{n-1} \phi_n$. To verify that Φ is bijective, Foata and Han introduce a class of bijections ψ_l , whose definitions are parallel to the definitions of the ϕ_l 's. Let $w = x_1 x_2 \cdots x_n \in \text{Sh}(0^{n-m}v)$ ($0 \leq n \leq m$). We define n bijections ψ_l ($1 \leq l \leq n$) of $\text{Sh}(0^{n-m}v)$ onto itself in the following manner: For each l such that $n - m + 1 \leq l \leq n$ let $\psi_l(w) = w$. When $1 \leq l \leq n - m$, let x_j denote the l -th letter of w , equal to 0, when w is read from left to right. Consider the following three cases (remember $x_0 = x_{n+1} = +\infty$ by convention):

- (1') x_{j-1}, x_{j+1} both non-subexcedent;
- (2') x_{j-1} subexcedent, x_{j+1} non-subexcedent or x_{j-1}, x_{j+1} both subexcedent with $x_{j-1} > x_{j+1}$;
- (3') x_{j-1} non-subexcedent, x_{j+1} subexcedent or x_{j-1}, x_{j+1} both subexcedent with $x_{j-1} < x_{j+1}$.

When case (1') holds, let $\psi_l(w) := w$. When case (2') holds, determine the smallest integer $k \leq j-1$ such that $x_k < \dots < x_{j-1} < \text{rank}(j-1)$ so that $w = x_1 \dots x_{k-1} x_k \dots x_{j-1} 0 x_{j+1} \dots x_n$ and define: $\psi_l(w) := x_1 \dots x_{i-1} 0 x_i \dots x_{j-1} x_{j+1} \dots x_n$. When case (3') holds, determine the greatest integer $k \geq j+1$ such that $\text{rank}(j+1) > x_{j+1} > x_{j+2} > \dots > x_k$ so that $w = x_1 \dots x_{j-1} 0 x_{j+1} \dots x_k x_{k+1} \dots x_n$ and define: $\psi_l(w) := x_1 \dots x_{j-1} x_{j+1} \dots x_k 0 x_{k+1} \dots x_n$. As shown in [7], the product $\Phi^{-1} := \psi_n \psi_{n-1} \dots \psi_1$ is the inverse bijection of Φ .

5.2. Foata-Han's second transformation F. The bijection $\mathbf{F}$ maps each shuffle class $\text{Sh}(0^{n-m}v)$ with v an arbitrary word of length m ($0 \leq m \leq n$) onto itself. When $n = 1$ the unique element of the shuffle class is sent onto itself. Also let $\mathbf{F}(w) = w$ when $\text{des}(w) = 0$. Let $n \geq 2$ and assume that $\mathbf{F}(w')$ has been defined for all words w' with nonnegative letters, of length $n' \leq n-1$. Let w be a word of length n such that $\text{des}(w) \geq 1$. We may write

$$w = w'a0^r b,$$

where $a \geq 1$, $b \geq 0$ and $r \geq 0$. Three cases are considered:

$$(1) \quad a \leq b; \quad (2) \quad a > b, \quad r \geq 1; \quad (3) \quad a > b, \quad r = 0.$$

In case (1) define $\mathbf{F}(w) = \mathbf{F}(w'a0^r b) := (\mathbf{F}(w'a0^r))b$. In case (2) we may write $\mathbf{F}(w'a0^r) = w''0$ and then define

$$\begin{aligned} \gamma \mathbf{F}(w'a0^r) &:= 0w''; \\ \mathbf{F}(w) = \mathbf{F}(w'a0^r b) &:= (\gamma \mathbf{F}(w'a0^r))b = 0w''b. \end{aligned}$$

In short, add one letter “0” to the left of $\mathbf{F}(w'a0^r)$, then delete the rightmost letter “0” and add b to the right. In case (3) remember that $r = 0$. Write

$$\mathbf{F}(w'a) = 0^{m_1} x_1 v_1 0^{m_2} x_2 v_2 \dots 0^{m_k} x_k v_k,$$

where $m_1 \geq 0$, $m_2, \dots, m_k$ are all positive, then $x_1, x_2, \dots, x_k$ are positive letters and $v_1, v_2, \dots, v_k$ are words with positive letters, possibly empty. Then define:

$$\begin{aligned} \delta \mathbf{F}(w'a) &:= x_1 0^{m_1} v_1 x_2 0^{m_2} v_2 x_3 \dots x_k 0^{m_k} v_k; \\ \mathbf{F}(w) = \mathbf{F}(w'ab) &:= (\delta \mathbf{F}(w'a))b. \end{aligned}$$

In short, move each positive letter occurring just after a 0-factor of $\mathbf{F}(w'a)$ to the beginning of that 0-factor and add b to the right.

5.3. Proof of Theorem 3. We show that the composition $\text{ZDer}^{-1} \circ \mathbf{F} \circ \Phi^{-1} \circ \text{ZDer}$ satisfies the relations (3.4)–(3.7) characterizing the bijection Ψ . The proof is based on Lemmas 3, 5 and 6, which will be proved in Section 6.

Let $\tau = \langle \sigma, i_1, i_2, \dots, i_m \rangle \in \mathcal{S}_{n+m}$, where $\sigma = x_1 \dots x_n \in \mathcal{D}_n$ is the derangement part of τ . Since the positions of fixed points of τ are the same as that of zeros of $\text{ZDer}\tau$, we write

$$\text{ZDer}\tau = [\sigma, i_1, i_2, \dots, i_m] \in \text{Sh}(0^m \sigma).$$

Thus the k -th zero from left to right of $\text{ZDer}\tau$ is at the position $i_k + k$ for $k \leq m$ and i_k is the number of pillars at the left of the k -th zeros in $\text{ZDer}\tau$. Consequently, writing $i := i_m$ then

$$[x_1 x_2 \dots x_n, i_1, i_2, \dots, i_m] = w x_i 0^r x_{i+1} x_{i+2} \dots x_n, \quad (5.1)$$

where w is a word with $i - 1$ pillars and r the largest integer satisfying $i_{m+1-r} = i$. The integer r must be positive for the last zero is located just at the left of the $(i + 1)$ -th pillar.

Example. Let $\Omega = [3\ 1\ 4\ 5\ 2, 0, 0, 0, 1, 1, 2, 2, 2, \mathbf{2}]$. Then $i = 2$, $r = 4$, $w = 000300$ and

$$\Omega = w\ 1\ 0^4\ 4\ 5\ 2 = 000300\mathbf{1}0000452.$$

For brevity we introduce the following notations: $\forall t \geq 0$ and $I = (i_1, \dots, i_m) \in \mathbb{N}^m$, let $I + t = (i_1 + t, \dots, i_m + t)$, $t^{[m]} = (t, \dots, t) \in \mathbb{N}^m$. Moreover, for any $\omega \in \text{Sh}(0^{n-m}v)$, if $\mathbf{F}(\omega) = [\sigma, i_1, \dots, i_m]$, then we write

$$\mathbf{F}(\omega) = [\sigma, I_\omega] \quad \text{with} \quad I_\omega = (i_1, \dots, i_m).$$

Additionally, for any non empty finite word w we denote, respectively, by $L(w)$ and $R(w)$ the first and last letter of w when w is read from left to right.

Lemma 3. *Let w_1, w_2 be two non empty words such that $R(w_1) > 0$ and $\text{zero}(w_2) = 0$. Assume that $(R(w_1), L(w_2)) = (a, b)$ and $\text{pil}(w_1) = \nu$, $\text{des}(w_2) = t$. Let $\mu = w_1 0^r w_2$ with $r \geq 0$. Then*

$$\text{zero}(w_1) \neq 0 \implies I_\mu = \begin{cases} (t^{[r]}, I_{w_1} + t + 1), & \text{if } a > b, \\ I_{w_1} + t, & \text{if } a < b \text{ and } r = 0, \\ (t^{[r-1]}, I_{w_1} + t + 1, \nu + t), & \text{if } a < b \text{ and } r > 0; \end{cases} \quad (5.2)$$

and

$$\text{zero}(w_1) = 0 \implies I_\mu = \begin{cases} \emptyset, & \text{if } r = 0, \\ (t^{[r]}), & \text{if } r > 0 \text{ and } a > b, \\ (t^{[r-1]}, \nu + t), & \text{if } r > 0 \text{ and } a < b. \end{cases} \quad (5.3)$$

Let $\sigma = x_1 x_2 \dots x_n \in S_n$. The color of a slot $(i, i + 1)$ of σ can be characterized (see [4]) as follows:

- The slot $(i, i + 1)$ of σ is green if and only if one of the following conditions is satisfied:

$$(G_1) \quad x_i > x_{i+1} > i; \quad (G_2) \quad x_i < i < x_{i+1}; \quad (G_3) \quad i > x_i > x_{i+1}. \quad (5.4)$$

- The slot $(i, i + 1)$ of σ is red if and only if one of the following conditions is satisfied:

$$(R_1) \quad i < x_i < x_{i+1}; \quad (R_2) \quad x_i > i \geq x_{i+1}; \quad (R_3) \quad x_i < x_{i+1} \leq i. \quad (5.5)$$

By convention $x_0 = x_{n+1} = +\infty$. If we denote by d_i the number of descents of $\langle \sigma, i \rangle$ at right of i ($\sigma \in \mathcal{D}_n$) then Lemma 1 implies that $g_i = d_i$ if the i -th slot is green, and $g_i = d_i + i$ if the slot is red.

Lemma 4. *Let $\sigma = x_1 x_2 \dots x_n \in S_n$, let $(i, i + 1)$ be the j -th slot of σ and $t_i = \text{des}(x_{i+1} \dots x_n)$ ($1 \leq i \leq n$) then*

$$d_j = \begin{cases} t_i & \text{in the cases } G_1, G_2, \text{ or } R_1; \\ t_i + 1 & \text{in the cases } G_3, R_2, \text{ or } R_3. \end{cases}$$

Proof. Clearly, in the cases G_1, G_2, R_1 the value x_{i+1} is an exceedant of σ , so $i + 1$ is not a descent place of $\langle \sigma, j \rangle$ while in the cases G_3, R_2 and R_3 the value x_{i+1} is a sub-exceedant, so $i + 1$ is a descent place of $\langle \sigma, j \rangle$. $\square$

Let $\tau = \langle \sigma, i_1, i_2, \dots, i_m \rangle \in S_{n+m}$ and $\Omega = \text{ZDer}\tau = [\sigma, i_1, \dots, i_m]$. We distinguish two cases according to the color of the insertion slot.

5.4. The i_m -th slot of σ is green. Define

$$\omega = [\sigma, i_1, \dots, i_{m-1}], \quad \omega' = \Phi^{-1}(\omega) \quad \text{and} \quad \Omega' = \Phi^{-1}(\Omega).$$

We must check (3.4) and for $m > 1$,

$$\mathbf{F} \circ \Phi^{-1}(\omega) = [\sigma, j_2, \dots, j_m] \implies \mathbf{F} \circ \Phi^{-1}(\Omega) = [\sigma, g_{i_m}, j_2, \dots, j_m], \quad (5.6)$$

which is equivalent to

$$I_{\Omega'} = (g_{i_m}, I_{\omega'}). \quad (5.7)$$

Lemma 5. *We have the following factorizations:*

$$\omega' = w_1 0^r w_2 0^{r'} w_3 \quad \text{and} \quad \Omega' = w_1 0^{r+1} w_2 0^{r'} w_3 \quad (r, r' \geq 0),$$

where $w_1 \neq \emptyset$, $w_2 \neq \emptyset$ and w_3 are words on non negative integers. Moreover, if $(a, b, a', b') = (\text{R}(w_1), \text{L}(w_2), \text{R}(w_2), \text{L}(w_3))$, then the following properties hold true:

- i) $\text{zero}(w_2 w_3) = 0$, and $\text{R}(w_1) > 0$,
- ii) if $r = 0$, then $a > b$,
- iii) if $r' > 0$, then $r' = 1$, $a' < b'$ and $r = 0$,
- iv) $\text{des}(w_2 0^{r'} w_3) = g_{i_m}$.

Notice that if $r' = 1$ then $w_3 \neq \emptyset$ by iii). Let $w'_2 = w_2 w_3$. If $m = 1$ then $\omega = \omega' = \sigma$ and $\Omega = [\sigma, i_1]$. Hence $\text{zero}(\omega) = 0$ and $r = r' = 0$. Thus $\omega' = w_1 w'_2$ and $\Omega' = w_1 0 w'_2$, where $a > b = \text{L}(w'_2)$ and $\text{des}(w'_2) = g_{i_1}$. It follows then from (5.3) that

$$I_{\Omega'_1} = (g_{i_1}). \quad (5.8)$$

We now prove (5.7) for $m > 1$.

1) If $r' = 0$ then

$$\omega' = w_1 0^r w'_2 \quad \text{and} \quad \Omega' = w_1 0^{r+1} w'_2, \quad (5.9)$$

where $\text{des}(w'_2) = t = g_{i_m}$ and $\text{zero}(w'_2) = 0$. As $m > 1$ we cannot have $\text{zero}(w_1) = 0$ and $r = 0$ simultaneously. It remains to verify the following two cases:

- i) $\text{zero}(w_1) \neq 0$, $a > b$ or $r > 0$ and $a < b$,
- ii) $\text{zero}(w_1) = 0$, $r > 0$ and $a \neq b$.

Applying Lemma 3 to either case yields

$$I_{\Omega'} = (t, I_{\omega'}) = (g_{i_m}, I_{\omega'}). \quad (5.10)$$

2) If $r' = 1$, then $a' < b'$, $r = 0$ and $a > b$. Hence

$$\omega' = v 0 w_3 \quad \text{and} \quad \Omega' = V 0 w_3,$$

where $v = w_1 w_2$ and $V = w_1 0 w_2$. Let $t_2 = \text{des}(w_2)$, $t_3 = \text{des}(w_3)$ and $t = \text{des}(w_2 0 w_3) = g_{i_m}$, $\nu = \text{pil}(v) = \text{pil}(V)$. If $\text{zero}(w_1) \neq 0$, as $a > b$, by (5.2) we have

$$I_v = I_{w_1} + t_2 + 1 \quad \text{and} \quad I_V = (t_2, I_{w_1} + t_2 + 1) = (t_2, I_v). \quad (5.11)$$

Since $\text{zero}(v) > 0$, $\text{zero}(V) > 0$ and $a' < b'$, by (5.2) we have

$$I_{\omega'} = (I_v + t_3 + 1, \nu + t_3) \quad \text{and} \quad I_{\Omega'} = (I_V + t_3 + 1, \nu + t_3). \quad (5.12)$$

It follows that $I_{\Omega'} = (t_2 + t_3 + 1, I_{\omega'})$ and (as $a' < b'$) $t = t_2 + t_3 + 1$. Now, if $\text{zero}(w_1) = 0$ then, by (5.3),

$$I_{\omega'} = (\nu + t_3) \quad \text{and} \quad I_V = (t_2). \quad (5.13)$$

As $\text{zero}(V) > 0$, it follows from (5.12) and (5.13) that $I_{\Omega'} = (t, \nu + t_3) = (t, I_{\omega'})$.

5.5. The i_m -th slot of σ is red. Let k be the largest integer such that $i_{m-k+1} = i_m$. Define

$$\omega = [\sigma, i_1, \dots, i_{m-k}], \quad \omega' = \Phi^{-1}(\omega), \quad \Omega' = \Phi^{-1}(\Omega) \quad \text{and} \quad \tilde{\Omega} = \psi_{m-k} \cdots \psi_1(\Omega).$$

For $k = m$ the relation (3.6) is equivalent to

$$I_{\Omega'} = ((g_{i_m} - i_m)^{[m-1]}, g_{i_m}), \quad (5.14)$$

which corresponds to (3.4) when $m = 1$. For $k < m$ the relation (3.7) is equivalent to

$$I_{\Omega'} = ((g_{i_m} - i_m)^{[k-1]}, I_{\omega'} + 1, g_{i_m}). \quad (5.15)$$

We now verify (5.14) and (5.15). Recall that $\sigma = x_1 \dots x_n$. For convenience, we use write $i := i_m$ in what follows.

5.5.1. Assume that x_{i+1} is excedent. This corresponds to the case (R_1) , i.e., x_i and x_{i+1} are both excedent and $x_i < x_{i+1}$. Hence, all the zeros of ω are at the left of x_i and all the zeros of ω' remains at the left of x_i . It follows that

$$\omega' = w_1 x_i x_{i+1} \cdots x_n \quad \text{and} \quad \tilde{\Omega} = w_1 x_i 0^k x_{i+1} \cdots x_n \quad (5.16)$$

Besides, as the map ψ_j is identity for $m - k + 1 \leq j \leq m$, we have $\Omega' = \tilde{\Omega}$. Let $t = \text{des}(x_{i+1} \cdots x_n)$. By Lemma 4 we have $t = g_i - i$.

If $\text{zero}(w_1) = 0$ then $m = k$. From (5.3) we derive $I_{\Omega'} = (t^{[k-1]}, i + t) = ((g_i - i)^{[m-1]}, g_i)$, which is (5.14). If $\text{zero}(w_1) \neq 0$, by (5.2) we have

$$I_{\omega'} = (I_{w_1 x_i} + t) \quad \text{and} \quad I_{\Omega'} = (t^{[k-1]}, I_{w_1 x_i} + t + 1, i + t),$$

which is precisely (5.15).

5.5.2. Assume that x_{i+1} is subexcedent. We need the following result.

Lemma 6. *Let h be the largest integer such that $i + 1 > x_{i+1} > \cdots > x_h$ and $\ell = i + 1$ if x_i is excedent or the last zero of w' is located between x_i and x_{i+1} otherwise ℓ be the smallest integer such that $x_\ell < \cdots < x_{i+1} < i + 1$ and that ω' does not contain zero at the right of x_ℓ . Set $T = \text{des}(x_\ell \cdots x_n)$; $w_3 = x_{h+1} \cdots x_n$; $t' = \text{des}(w_3)$, then we have*

$$\omega' = w_1 x_{\ell-1} 0^r x_\ell \cdots x_h w_3, \quad (5.17)$$

$$\Omega' = w_1 x_{\ell-1} 0^{r+k-1} x_\ell \cdots x_h 0 w_3. \quad (5.18)$$

Moreover, the following identities hold true:

- i) $(r = 0 \text{ and } x_{\ell-1} > x_\ell) \text{ or } (r = 1 \text{ and } x_{\ell-1} < x_\ell),$
- ii) $T + 1 = g_i - i,$
- iii) $h + t' = g_i.$

Let $t = \text{des}(x_\ell \cdots x_h)$. As $x_h < x_{h+1}$ we have $T = t + t'$. Write $\Omega' = V0w_3$ with

$$V = w_1 x_{\ell-1} 0^{r+k-1} x_\ell \cdots x_h.$$

- Suppose that $k = m$. Then $r = 0$, $\text{zero}(w_1 x_{\ell-1}) = 0$ and $x_{\ell-1} > x_\ell$.
 - If $k = 1$ then $\text{zero}(V) = 0$. By (5.3) we have $I_{\Omega'} = (h + t') = (g_i)$, which is equivalent to (3.4).
 - If $k > 1$ then $\text{zero}(V) > 0$. By (5.2) we have $I_{\Omega'} = (I_V + t' + 1, h + t')$, while (5.3) yields $I_V = t^{[k-1]}$. Hence $I_{\Omega'} = ((T + 1)^{[k-1]}, h + t')$, which is equivalent to (5.14).
- Suppose that $k < m$. There are three cases:
 - (a) $r = 0$, we have $\text{zero}(w_1) \neq 0$, $x_{\ell-1} > x_\ell$ and, by (5.2),

$$I_{\omega'} = (I_{w_1 x_{\ell-1}} + T + 1) \quad \text{and} \quad I_V = (t^{[k-1]}, I_{w_1 x_{\ell-1}} + t + 1).$$

- (b) $r = 1$ and $\text{zero}(w_1) \neq 0$, we have $x_{\ell-1} < x_\ell$. By (5.2)

$$I_{\omega'} = (I_{w_1 x_{\ell-1}} + T + 1, \ell - 1 + T) \quad \text{and} \quad I_V = (t^{[k-1]}, I_{w_1 x_{\ell-1}} + t + 1, \ell - 1 + t).$$

- (c) $r = 1$ and $\text{zero}(w_1) = 0$, by (5.3) we have

$$I_{\omega'} = (\ell - 1 + T) \quad \text{and} \quad I_V = (t^{[k-1]}, \ell - 1 + t).$$

On the other hand, in any case, we have $\text{zero}(V) > 0$. By (5.2),

$$I_{\Omega'} = (I_V + t' + 1, h + t') = ((T + 1)^{[k-1]}, I_{\omega'} + 1, h + t'),$$

which is equivalent to (5.15).

Example. Let $\tau = 15342768 \in S_8$ then $\tau = \langle \sigma, 0, 1, 1, 4 \rangle$, where $\text{Der} \tau = \sigma = 2143$. So $\langle \sigma, 0 \rangle = 13254$; $\langle \sigma, 1 \rangle = 32154$; $\langle \sigma, 2 \rangle = 21354$; $\langle \sigma, 3 \rangle = 21543$. $\langle \sigma, 4 \rangle = 21435$; hence the slots 0, 2, 4 are green, while the slots 1 and 3 are red. Therefore $(g_0, \dots, g_4) = (2, 3, 1, 4, 0)$, and

- slot 0 is green $\implies \Psi \langle \sigma, 0 \rangle = \langle \sigma, g_0 \rangle = \langle \sigma, 2 \rangle$,
- slot 1 is red $\implies \Psi \langle \sigma, 0, 1, 1 \rangle = \langle \sigma, g_1 - 1, 2 + 1, g_1 \rangle = \langle \sigma, 2, 3, 3 \rangle$,
- slot 4 is green $\implies \Psi \langle \sigma, 0, 1, 1, 4 \rangle = \langle \sigma, g_4, 2, 3, 3 \rangle = \langle \sigma, 0, 2, 3, 3 \rangle$.

Thus $\Psi(\tau) = \langle \sigma, 0, 2, 3, 3 \rangle = 13248675$.

On the other hand, applying Φ^{-1} to $w := \text{ZDer} \tau = 02001430 \in \text{Sh}(0^4 \sigma)$, as 2 and 4 are excedent, we get

$$\begin{array}{rcll}
 Id & & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 \\
 w & = & 0 & 2 & 0 & 0 & 1 & 4 & 3 & 0 \\
 \psi_1(w) & = & 0 & 2 & 0 & 0 & 1 & 4 & 3 & 0 & \text{Case (1')} \\
 \psi_2(\psi_1(w)) & = & 0 & 2 & 0 & 0 & 1 & 4 & 3 & 0 & \text{Case (1')} \\
 \psi_3(\psi_2(\psi_1(w))) & = & 0 & 2 & 0 & 1 & 0 & 4 & 3 & 0 & \text{Case (3')} \text{ with } k = 5 \\
 \psi_4(\psi_3(\psi_2(\psi_1(w)))) & = & 0 & 2 & 0 & 1 & 0 & 4 & 0 & 3 & \text{Case (2')} \text{ with } k = 7.
 \end{array}$$

Thus $\Phi^{-1}(w) = 02010403$. Now we apply $\mathbf{F}$ to $w = 02010403$:

$\mathbf{F}(02) = 02$	no descent
$\mathbf{F}(020) = \delta(02)0 = 200$	Case (3)
$\mathbf{F}(0201) = \gamma(200)1 = 0201$	Case (2)
$\mathbf{F}(02010) = \delta(0201)0 = 20100$	Case (3)
$\mathbf{F}(020104) = 201004$	Case (1)
$\mathbf{F}(0201040) = \delta(020104)0 = 2104000$	Case (3)
$\mathbf{F}(02010403) = \gamma(2104000)3 = 02104003$	Case (2).

So $\mathbf{F}(02010403) = [\sigma, 0, 2, 3, 3]$ and $\text{ZDer}^{-1}([\sigma, 0, 2, 3, 3]) = \langle \sigma, 0, 2, 3, 3 \rangle = 1\,3\,2\,4\,8\,6\,7\,5$. Therefore (2.10) is checked.

6. THE PROOF OF THREE LEMMAS

6.1. Proof of Lemma 3. Recall that δ and Υ are the transformations used in the cases (b) and (c) of the algorithm $\mathbf{F}$. Let α a word of length ν on the alphabet of positive integers and $b > 0$. Let $w = [\alpha, i_1, \dots, i_z] \in \text{Sh}(0^z\alpha)$. Note that

$$wb = [\alpha b, i_1, \dots, i_z], \quad w0 = [\alpha, i_1, \dots, i_z, \nu], \quad 0w = [\alpha, 0, i_1, \dots, i_z].$$

Hence, if $\text{R}(w) = 0$, i.e., $i_z = \nu$, then $\Upsilon[\alpha, i_1, \dots, i_z] = [\alpha, 0, i_1, \dots, i_{z-1}]$. On the other hand, if $x_1, \dots, x_n$ are positive integers, then

$$x_1 \cdots x_k 0^r x_{k+1} \cdots x_n = [x_1 \cdots x_n, k^{[r]}].$$

It follows that $\delta(w) = [\alpha, i_1 + 1, \dots, i_z + 1]$. Now, consider the word $\mu = w_1 0^r w_2$, where $\text{Pil}(\mu) = \sigma$, $\text{Pil}(w_1) = \alpha$ and $|\alpha| = \nu$. Since w_1 and w_2 are non empty we can write $w_1 = v_1 a$ and $w_2 = b v_2$. Therefore $\mu = v_1 a 0^r b v_2$.

If $\text{zero}(w_1) = z \neq 0$ and $a > b$ set $I_{w_1} = (\ell_1, \dots, \ell_z)$. Then, we have successively

$$\mathbf{F}(v_1 a 0) = \delta(\mathbf{F}(v_1 a))0 = [\alpha, \ell_1 + 1, \dots, \ell_z + 1, \nu] \implies I_{v_1 a 0} = (I_{v_1 a} + 1, \nu),$$

$$\mathbf{F}(v_1 a 0 0) = \Upsilon(\mathbf{F}(v_1 a 0))0 = [\alpha, 0, \ell_1 + 1, \dots, \ell_z + 1, \nu] \implies I_{v_1 a 0^r} = (0^{[r-1]}, I_{v_1 a} + 1, \nu),$$

and $\mathbf{F}(v_1 a 0^r b) = \Upsilon(\mathbf{F}(v_1 a 0^{[r]}))b = [\alpha b, 0^r, I_{w_1} + 1]$. Finally, as $\text{zero}(v_2) = 0$, we have $\mathbf{F}(\mu) = [\sigma, t^{[r]}, I_{w_1} + 1 + t]$, which corresponds to the first case of (5.2). The other cases can be proved similarly.

6.2. Proof of Lemma 5. Recall that $\Omega = [\sigma, i_1, \dots, i_m]$, where

$$\sigma = x_0 x_1 \cdots x_n x_{n+1}, \quad \omega = [\sigma, i_1, \dots, i_{m-1}], \quad \Omega' = \Phi^{-1}(\Omega), \quad \omega' = \Phi^{-1}(\omega).$$

Set $\tilde{\omega} = \psi_{m-2} \circ \cdots \circ \psi_1(\omega)$, $\tilde{\Omega} = \psi_{m-1} \circ \cdots \circ \psi_1(\Omega)$ and $i = i_m$. The last zero of ω is at the left of x_{i+1} so is the last zero of $\tilde{\omega}$ by definition of Φ^{-1} . So all $m - 2$ zeros of ω' are on the left of x_{i+1} and only the last can be on the left or on the right of x_{i+1} . Hence ω' is of the following form:

$$\omega' = v_1 x_i 0^\epsilon x_{i+1} x_{n+2} \cdots x_h 0^{r'} x_{h+1} \cdots x_n x_{n+1},$$

where $\epsilon \geq 0$, and $0 \leq r' \leq 1$. Keeping in mind the definition of Φ^{-1} , if x_{i+1} is excedent (the case of G_1 or G_2) then we have necessarily $r' = 0$ and If x_{i+1} is subexcedent (the case of G_3)

then it is impossible for the last zero of ω' to be between x_i and x_{i+1} , otherwise when one applies to ω' the map ψ_{m-1}^{-1} , it corresponds to the case (2) therefore the last zero of $\tilde{\omega}$ would be on the right of x_{i+1} , that is impossible. So in case of G_3 we have necessarily $r' = 1$ or ($r' = 0$ and $\epsilon = 0$). We consider the following three cases:

- G_1 or (G_2 and $\epsilon > 0$). Since $r' = 0$, hence ω' and $\tilde{\omega}$ are of the following forms:

$$\omega' = v_1 x_i 0^\epsilon x_{i+1} \dots x_{n+1}, \quad \tilde{\Omega} = v_1 x_i 0^{\epsilon+1} x_{i+1} \dots x_{n+1}.$$

In this case ψ_m is the identity therefore $\Omega' = \tilde{\Omega}$ and then we have

$$r = \epsilon; \quad r' = 0 \quad w_1 = v_1 x_i; \quad w_2 = x_{i+1} \dots x_n x_{n+1}; \quad w_3 = \emptyset.$$

If $r = 0$ we have G_1 so $a > b$ and by Lemma 4, $\text{des}(w_2) = g_i$.

- (G_2 and $\epsilon = 0$) or (G_3 and $r' = 0$). In this case, let k be the smallest integer such that $x_k < \dots < x_i < i$ and ω' does not contain any zero on the right of x_k . Thus ω' and $\tilde{\Omega}$ are of the following forms:

$$\begin{aligned} \omega' &= v_1 x_{k-1} 0^r x_k \dots x_i x_{i+1} \dots x_{n+1}, \\ \tilde{\Omega} &= v_1 x_{k-1} 0^r x_k \dots x_i 0 x_{i+1} \dots x_{n+1}. \end{aligned}$$

Since $\Omega' = \psi_m(\tilde{\Omega})$ and the map ψ_m corresponds to the case (2'), we have

$$\Omega' = v_1 x_{k-1} 0^{r+1} x_k \dots x_i x_{i+1} \dots x_n x_{n+1}.$$

Moreover $r' = 0$ $w_1 = v_1 x_{k-1}$ and $w_2 = x_k \dots x_n x_{n+1}$; $w_3 = \emptyset$. As $\text{des}(x_k \dots x_i) = 0$, we have $\text{des}(w_2) = g_i$ by Lemma 4.

- (G_3 and $r' = 1$). In this case, both x_i and x_{i+1} must be subexcedances and $x_i > x_{i+1}$ and h must be the largest integer such that $i + 1 > x_{i+1} > \dots > x_h$ (by applying ϕ_{m-1} to $\tilde{\omega}$). Let k be the smallest integer such that $k > x_k > \dots > x_{i+1}$ and ω' contains only one zero on the right of x_k . Thus we have

$$\omega' = v_1 x_{k-1} 0^\alpha x_k \dots x_i x_{i+1} \dots x_h 0 x_{h+1} \dots x_{n+1}.$$

On the other hand, the last zero of $\tilde{\omega}$ is at the left of x_{i+1} , more precisely just at the left of x_k because $k > x_k > x_i > x_{i+1} > \dots > x_h$ by the map ϕ_{m-1} . Hence

$$\tilde{\omega} = v_1 x_{k-1} 0^{\alpha+1} x_k \dots x_i x_{i+1} \dots x_h x_{h+1} \dots x_{n+1}.$$

Set $\tilde{\tilde{\Omega}} = \psi_{m-2} \circ \dots \circ \psi_1(\tilde{\Omega})$. Then

$$\tilde{\tilde{\Omega}} = [\tilde{\omega}, i] = v_1 x_{k-1} 0^{\alpha+1} x_k \dots x_i 0 x_{i+1} \dots x_h x_{h+1} \dots x_{n+1}.$$

When we apply ψ_{m-1} to $\tilde{\tilde{\Omega}}$, it corresponds to the case (3'), so

$$\tilde{\tilde{\Omega}} = \psi_{m-1}(\tilde{\tilde{\Omega}}) = v_1 x_{k-1} 0^\alpha x_k \dots x_i 0 0 x_{i+1} \dots x_h x_{h+1} \dots x_{n+1}.$$

Similarly we have

$$\Omega' = \psi_m(\tilde{\tilde{\Omega}}) = v x_{k-1} 0^\alpha x_k \dots x_i 0 x_{i+1} \dots x_h 0 x_{h+1} \dots x_{n+1}.$$

In this situation we take $r = 0$ ($r' = 1$) and

$$w_1 = v_1 x_{k-1} 0^\alpha x_k \dots x_i; \quad w_2 = x_{i+1} \dots x_h; \quad w_3 = x_{h+1} \dots x_{n+1}.$$

As $x_h < x_{h+1}$ we have $\text{des}(w_2 0 w_3) = \text{des}(w_2 w_3) + 1 = g_i$ by Lemma 4.

6.3. Proof of Lemma 6. Since the i -th slot of σ is red and x_{i+1} is subexcedent, we are in the situation of R_2 or R_3 . Recall that all the *zeros* of ω are located at the left of x_i so are all of $\tilde{\omega} = \psi_{m-k-1} \circ \cdots \circ \psi_1(\omega)$. We show that the last zero of ω' is at the left of x_{i+1} . In the R_2 case x_i is excedent and all zeros of ω' are on the left of x_i . In the R_3 case, suppose that the last zero of ω' is at the right of x_{i+1} , then, by applying the reverse mapping $\psi_{m-k}^{-1} = \phi_{m-k}$, the last *zero* of $\tilde{\omega}$ cannot be at the left of x_i because $x_i < x_{i+1}$. This is absurd. In order to show (5.17) and (5.18) set $\Omega_{m-k} = \tilde{\Omega}$ and for all j such that $m-k < j \leq m$ set $\Omega_j = \psi_j(\Omega_{j-1})$. There are two cases:

- If $\ell \leq i$ then

$$\omega' = w_1 x_{\ell-1} 0^r x_\ell \cdots x_i x_{i+1} \cdots x_h x_{h+1} w_3,$$

and

$$\tilde{\Omega} = w_1 x_{\ell-1} 0^r x_\ell \cdots x_i 0^k x_{i+1} \cdots x_h x_{h+1} w_3.$$

Noticing that $x_i < x_{i+1} < i+1$, so x_i is subexcedent and the application of ψ_j to Ω_{j-1} corresponds to case (2') for all $m-k < j < m$ and ψ_m corresponds to case (3'). Therefore $\Omega' = w_1 x_{\ell-1} 0^{r+k-1} x_\ell \cdots x_i x_{i+1} \cdots x_h 0 x_{h+1} w_3$.

- If $\ell = i+1$, then $\omega' = w_1 x_i 0^r x_{i+1} \cdots x_h x_{h+1} w_3$ and $\tilde{\Omega} = w_1 x_i 0^{r+k} x_{i+1} \cdots x_h x_{h+1} w_3$. In this case, ψ_j corresponds to the case (1) for all $m-k < j < m$, and ψ_m corresponds to case (3'). So $\Omega' = w_1 x_i 0^{r+k-1} x_{i+1} \cdots x_h 0 x_{h+1} w_3$.

It remains to verify the three conditions of Lemma 6 in the above two cases. It is clear that $x_{\ell-1} > x_\ell$ if $r = 0$. Moreover, neither $r > 1$ nor ($r = 1$ and $x_{\ell-1} > x_\ell$) is possible because, otherwise, when we apply ϕ_{m-k} to ω' , it corresponds to case (2), so the last *zero* of $\tilde{\omega}$ would be at the right of x_{i+1} , but this is absurd. So the condition (i) is verified. Besides, as $\text{des}(x_\ell \cdots x_{i+1}) = 0$ we have $T = \text{des}(x_{i+1} \cdots x_n)$, and by Lemma 4, we derive the condition (ii). Finally, by definition of h we have $\text{des}(x_{i+1} \cdots x_h) = h - i - 1$, and $T = \text{des}(x_{i+1} \cdots x_n) = h - i - 1 + t'$. Thus $h + t' = g_i$.

Acknowledgments: This work was done during the first author's visit to Institut Camille Jordan, Université Lyon 1 in the fall of 2008 and was supported by a scholarship of Agence universitaire de la francophonie. The second author acknowledges the financial support from la Région Rhône-Alpes via the program MIRA recherche 2008, projet: 0803414701.

REFERENCES

- [1] G. E. Andrews, *The theory of partitions*, Cambridge Mathematical Library, Cambridge University Press, Cambridge, 1998, Reprint of the 1976 original.
- [2] R. M. Adin, F. Brenti and Y. Roichman, Descent numbers and major indices for the hyperoctahedral group, *Adv. in Appl. Math.* **27** (2001), 210–224.
- [3] R. M. Adin and Y. Roichman, The flag major index and group actions on polynomial rings, *European J. Combin.* **22** (2001), 431–446.
- [4] R. Clarke, G.N. Han and J. Zeng, A combinatorial interpretation of the Seidel generation of q -derangement numbers, *Ann. Comb.* **1** (1997), no. 4, 313–327.
- [5] C. Chow, On derangement polynomials of color B , *Sém. Lothar. Combin.* **55** (2006), Art. B55b, 6 pp. (electronic).

- [6] H. Faliharimalala and J. Zeng, Derangements and Euler's difference table for $C_\ell \wr S_n$, *Electronic J. Combinatorics*, R65 of Volume 15(1), 2008.
- [7] D. Foata and G.-N. Han, Fix-Mahonian Calculus, I: two transformations, *Europ. J. Combinatorics*, Volume 29, Issue 7(2008), 1721-1732.
- [8] I. Gessel and C. Reutenauer, Counting permutations with given cycle structure and descent set, *J. Combin. Theory Ser. A*, vol. 64, 1993, p. 189–215.
- [9] M. Geck and G. Malle, Reflection groups, in: *Handbook of Algebra*, vol. 4 (edited by M. Hazewinkel), Elsevier, Amsterdam, 2006, 337-383.
- [10] J. Haglund, N. Loehr and J. B. Remmel, Statistics on wreath products, perfect matchings and signed words, *European J. Combin.* **26** (2005), 835–868.
- [11] V. Reiner, Signed permutation statistics, *European J. Combin.* 14 (1993), no. 6, 553–567.
- [12] R. P. Stanley, *Enumerative Combinatorics*, vol. 1, Cambridge University Press, Cambridge, 1997.
- [13] E. Steingr msson, Permutation statistics of indexed permutations, *European J. Combin.* 15 (1994), no. 2, 187–205.
- [14] M. Wachs, On q -derangement numbers, *Proc. Amer. Math. Soc.* **106** (1989), 273–278.
- [15] J. Zeng, The Akiyama-Tanigawa algorithm for Carlitz's q -Bernoulli numbers, *Integers: Electronic J. Combin. Number Theory* 6 (2006), #05.

D  PARTEMENT DE MATH  MATIQUES ET INFORMATIQUE, UNIVERSIT   D'ANTANANARIVO, 101 ANTANANARIVO, MADAGASCAR

E-mail address: heritianamihanta@yahoo.fr

UNIVERSIT   DE LYON, UNIVERSIT   LYON 1, INSTITUT CAMILLE JORDAN, UMR 5208 DU CNRED, F-69622, VILLEURBANNE CEDEX, FRANCE

E-mail address: zeng@math.univ-lyon1.fr